



*Annales Universitatis Paedagogicae Cracoviensis*

*Studia Naturae*, 9: XX–XX, 2024

ISSN 2543-8832

DOI: 10.24917/25438832.9.x

---

Mohamad Hesam Shahrajabian

Department of Agronomy and Plant Breeding, Isfahan (Khorasgan) Branch, Islamic Azad University, Isfahan 81551-39998, Iran; Correspondence: hesamshahrajabian@gmail.com

## Effects of climate change on functional health foods and medicinal plants – a short review

### Abstract

There is a growing opportunity to initiate studies to address the effects of climate change on medicinal plants' phenology, habitat alteration, species range shifts, and secondary metabolite production. Awareness of the potential effects of warming (increased CO<sub>2</sub>, and ultraviolet radiation due to ozone layer depletion) on secondary plant components and metabolites is an influential task for the future. There is information that climate change is causing remarkable effects on life cycles and the distribution of plant species. They are also forcing ecosystems to adopt the changing life cycle of plants and the development of new physical traits. A quickly changing climate might benefit species that can extend their ranges rapidly or that can tolerate a wide range of climatic conditions, both traits shared by various invasive plants taxon. This review provides a summary of the impact of climate change on medicinal and functional health food plants.

**Keywords:** climatic factors, global warming, plant yields, species ranges

**Received:** [2024.10.05]

**Accepted:** [2024.11.13]

### Introduction

Climate change is one of the most widespread global environmental changes, which are threatening the biodiversity and functioning of terrestrial ecosystems (Bates et al., 2019; Yin et al., 2019). Three main components define it – global warming, changes in precipitation patterns, and increased occurrence of extreme weather events (Ruszkiewicz et al., 2019).

Climatic parameters including natural disasters are projected to alter and directly influence

fishery, and livestock and plant production in multiple ways (Ayllon et al., 2019). Changes in global climate variables have already significantly affected the growth and productivity of plants. Crops are vulnerable to extreme events caused by climate change, especially drought, which is already having negative consequences for food production and food insecurity (Katsini et al., 2022). Heat stress, excessive CO<sub>2</sub> concentrations, and fluctuating precipitation patterns have changed the plant nutrient interactions (Lemes et al., 2022). Its negatively impacts plant nutrition, which serves as a basic hurdle in the production of enough nutritious food to meet the demands of the growing global population (Shahzad, Rouached, 2022). In addition, these phenomenons can increase among others: occurrence and virulence of food-borne pathogens, increase in food-borne infections and poisoning, risk of food contamination with chemical factors.

Climate variability associated with climate change influences agricultural production and rural livelihoods throughout the world, and climate services address this issue by making new information products available to agricultural technicians, policymakers, and farmers in order to enhance their adaptive capability (Green et al., 2022; Raiten, Bremer, 2023). Ensuring food safety under a changing climate requires adaptation techniques (Duchenne-Moutien, Neetoo, 2021). Managing climate change effects is a global concern that needs local action, and the ability of governments and city authorities to start and implement interventions remains a potential pathway toward decreasing climate change effects (Cobbinah et al., 2019). The promotion of climate agriculture has an important propensity to improve the adaptive capacity of smallholder crop farmers.

This paper aims to consider the effects of climate change on functional foods, functional health foods and medicinal plants. The screening of articles process was done in three stages, namely screening of title, abstract, and full text. The information in this article was provided from randomised control experiments, review articles, and analytical studies and observations which were gathered from numerous literature sources such as Scopus, Google Scholar, PubMed, and Science Direct.

### Functional health foods and medicinal plants

Many diseases can be prevented by appropriate functional health foods and diet which can optimize health improvement capabilities (Yeung et al., 2018). The body systems influenced by nutritionally related disorders are nervous, reproductive, musculoskeletal, respiratory, gastrointestinal,

cardiovascular, skin, endocrine, mental health, and special senses (Choudhary, Tandon, 2009). A suitable diet culminates in a healthy, properly functioning gastrointestinal (GI) tract, leading to the attainment of healthy living, and proper human physiology (Cencis, Chingwaru, 2010). A healthy and balanced diet is therefore a very good complement to the prevention and treatment of various diseases.


According to the classification of health foods, two main classes are food with specific health functions and nutritional supplements. Food with health functions is categorised into three groups, such as modern health food, traditional health food, and functional health food (Supachaturat et al., 2017). A functional health food can be a natural product that may have significant biological components, or a food achieved via a technological intervention that may boost its level of biologically active ingredients (Butnariu, Sarac, 2019). The different definitions of functional food categories are presented in table 1.

**Tab. 1.** Global definition of functional food groups (according to Del Castillo et al., 2018)

<b>Global definition for functional food category</b>	<b>Popular functional food ingredients worldwide</b>
1. Natural food with increased composition by employing particular agronomic conditions.	1. Probiotics, prebiotics and synbiotics.
2. Food including health-promoting constituents.	2. Dietary fiber.
3. Food from which a component has been removed to produce less negative impacts on health.	3. Omega 3 fatty acids, oleic acids and phytosterols.
4. Food in which the nature of one or more of its components has been chemically increased for obtaining health benefits.	4. Phytoestrogens.
5. Food in which the bioavailability of one or more of its components has been boosted to improve the assimilation of the health promoting components.	5. Phenolic compounds.

Some of the substantial functional food constituents of plants as a kind of nutraceutical are carotenoids, flavonoids, and dietary fibers (Shandilya, Sharma, 2017). Common carbohydrates, chlorophyll, and proteins are categorised as primary components where as phenolics, terpenoids, and alkaloids are grouped into secondary components (Sharma et al., 2021) – Fig. 1. For example, Kotilainen et al. (2006) reported that different kinds of functional food are fortified products (like fruit juices fortified with vitamin C or margarine with plant sterol ester), probiotics and prebiotics, altered products (such as fibers as fat releasers in meat or ice cream products), and enhanced commodities (like eggs with increased omega-3 content obtained by altered chicken feed). On the other hand, Bayir et al. (2019) considered polyphenols as an essential functional food, due to their diverse chemical structure. The main classes of polyphenols are flavanoids, stilbenes, lignans, and

phenolic acids.



**Fig. 1.** The most significant sources of nutraceutical substances in functional foods (based on: Kotilainen et al., 2006; Ranga Rao, Ravishankar, 2018)

It is often believed that healthy functional foods are a type of medicinal food (Das et al., 2016). This may be justified by the fact that some species of common vegetable plants have medicinal properties and their herbal raw materials are commonly used in herbal medicine. Good examples include onions, garlic, and various species of algae used as an addition to dishes. Of course, the amount of herbal raw material used and the method of its preparation plays a very important role in achieving a health-promoting effect. If we have isolated specific medicinal substances in high concentrations, it is difficult to talk about functional food in this case, but rather we are already dealing with a medicine. The method of application of this type of substance is also completely different. Medicines are taken differently, and functional or medicinal foods are prepared and consumed in various method. For example, algal powders and extracts are used in the nutraceutical, food, and pharmaceutical industries as tablets, crystals, capsules, gels, and dietary supplements (Ranga Rao, Ravishankar, 2018) – Tab. 2.

**Tab. 2.** Utilisation of algae for food and health applications (according to Ranga Rao, Ravishankar, 2018)

<b>Algal species</b>	<b>Applications</b>
<i>Spirulina platensis</i> (= <i>Arthrospira platensis</i> Gomont)	Phycocyanin, phycoerythrin, and biomass for health

<i>Chlorella vulgaris</i> Beijerinck; <i>Chlorella</i> spp.	food, feed, pharmaceuticals, and cosmetics. Polysaccharides for dietary supplements, extracts for cosmetics; Biomass for health food and feed.
<i>Dunaliella salina</i> (Dunal) Teodoresco	$\beta$ -carotene for health food, feed, dietary supplements, and cosmetics.
<i>Haematococcus pluvialis</i> Flotow	Astaxanthin for health food, pharmaceuticals and feed additives.
<i>Chlamydomonas reinhardtii</i> P.A.Dangeard	Biomass for animal health and feed; environmental monitoring bioremediation, production of recombinant proteins.
<i>Isochrysis galbana</i> Parke	Fatty acids for animal nutrition
<i>Nannochloropsis oculata</i> (Droop) D.J.Hibberd	Lipids and fatty acids for animal nutrition; extracts for cosmetics
<i>Porphyra</i> spp.	Biomass for feed, food; extracts for cosmetics
<i>Porphyridium</i> spp.	Polysaccharides for nutrition, pharmaceuticals, and cosmetics, phycoerythrin and phycocyanin for pharmaceuticals, food, and cosmetics.
<i>Phaeodactylum tricornutum</i> Bohlin	Lipids and fatty acids for nutrition

On the other hand, leafy thallus of species of the genus *Porphyra* is used for direct consumption in unprocessed form. Often, therapeutic therapies based on chemical drugs are supplemented with an appropriate functional or therapeutic diet, selected depending on the type of disease. Comparing functional foods with drugs and medicinal foods is shown in table 3.

**Tab. 3.** Comparing functional foods with medical foods and drugs (according to Das et al., 2016)

Difference	Functional foods	Medical foods	Prescription drugs
Uses	Energy enhancement, bolster gut, disease risk reduction, weight management, bone or heart health, memory improvement.	Dietary management of a disease or condition with distinctive nutritional requirements.	Treatment of disease, symptom or condition
Method of obtainment	No prescription or supervision is needed, consumer selects	Used with medical supervision	Prescribed by health provider
Distribution channels	Supermarkets, drugstores, online, major retailers	Hospitals, drugstores, pharmacies, online	Hospitals, Pharmacies
Amount consumed	As desired	As needed	As prescribed

However, it is worth being aware that combining medications, supplements, and a health-promoting diet should always be consulted with specialists in conventional medicine.

## Climate change and functional foods

### *Symptoms and effects*

The influences of climate on ecosystems are complex, complicated, and difficult to predict, and

ecosystem responses to climate and other contemporary changes will reverberate through the whole biomes (Phillips, 2019). Climate-changing parameters such as the availability of ozone, heat stress, and high levels of atmospheric carbon dioxide perturb redox balance and have a significant impact on the photosynthetic metabolism of plants and environmental stress reactions (Kumar et al., 2021). Rhizosphere microbiome can stimulate root nutrient absorption, arranging crop immune response, and increasing stress resistance. However, climate changes have a very significant impact on these all parameters (Li et al., 2021).

There are several pathways through which climate-related characteristics may influence food safety (Tab. 4).

**Tab. 4.** Main effects of climate change on the reduction of agricultural food production, both plant and animal origin (according to Semba et al., 2022)

Climate changes	<ol style="list-style-type: none"> <li>1) Extreme weather like drought, storms, heat waves;</li> <li>2) Ocean hypoxia; variations in sea level (flooding);</li> <li>3) Changes in temperature;</li> <li>4) Increase atmospheric carbon dioxide;</li> <li>5) Loss of coastal land, reefs, mangroves;</li> <li>6) Increase land degradation;</li> <li>7) Increase ground level ozone;</li> <li>8) Increase level of biotic stressors, decrease numbers of pollinators, and increase pathogens, and others.</li> </ol>
-----------------	---

These include, for example: increased repetition and intensity of extreme weather events, variations in temperature and precipitation patterns, acidification and ocean warming, and changes in the transport pathways of complicated contaminants, as already mentioned in the introduction (Ponce, 2020; Sun et al., 2023). Zhang and Liu (2022) discovered that climate changes prolonged the length of vegetative growth period, and they alter plant phenology. For example, alpine plants advances in the timing of first flowering alone will not necessarily translate to increases in flowering duration and enhanced reproductive effort. Instead, the findings demonstrate the importance of the date of last flowering in plant reproductive effort (Dorji et al., 2020). On the other hand, Wang et al. (2023) reported that the growth indicators of both C3 and C4 crops were more strongly associated with temperature and solar radiation, and variations in temperature and solar radiation had adverse effects on both C3 and C4 crops in different regions (Raes et al., 2021).

While warming can potentially reduce the crop growth period, water productivity and crop production are expected to decline because of higher water requirement and lower precipitation under higher temperature (Gohari et al., 2013). A marked increase in temperature since the middle

of the last century most probably influenced soil P dynamics with a significant effect on crop production (Seidel et al., 2021). According to many explorers, the leading cause of crop yield reductions is related to the soil water stress resulting from increased temperature and evapotranspiration during the growing season (Guo et al., 2021; Kang et al., 2022). The results on investigations show a frequent increase in the duration of dry season (an earlier onset of drought) and higher dependence on shallow precipitation water (Rolo, Moreno, 2019; Sun et al., 2024).


Among the important consequences of climate change is also worth mentioning the seasonal distribution of rain erosivity, which is a consequence of heavy downpours (Auerswald, Menzel, 2021). These factors can e.g. influence cover crop performance and soil nitrogen leaching events concomitantly (Teixeira et al., 2021). It is changing also the distributions and effects of plant diseases and pests. Climate warming favors the emergence of fungal pathogens, which can pose a significant threat to agricultural crops. Such phenomena concern also crop pests, the ranges of which are subject to shifts due to warming. Both of these factors have a very significant impact on the yield of functional food crops (Dorji et al., 2020; Raza, Bebbber, 2022).

In addition to all the important effects described above that affect the quantity of plant functional foods produced, it is also necessary to mention those factors resulting from climate change that affect the quality of the food produced. Several authors have noted that with climate change, microorganisms in the food supply chain have numerous negative effects (e.g. Alvi et al., 2021; Weber et al., 2023; Zhang et al., 2023). An increase in average temperatures could induce an increased risk of the proliferation of micro-organisms, like *Campylobacter* sp. and *Salmonella* sp., that produce food-borne illnesses (Miron et al., 2023). They are most often the cause of food poisoning, which in extreme cases can lead to death.

### *Remediation programs*

Considering crop growth, and crop disease models to predict the effects of climate change we can promote climate change adaptation techniques to ensure future food security (Newbery et al., 2016)

– Fig. 2.



**Fig. 2.** Integration of various adaptation/mitigation techniques and their potential positive effects to cope up with future and current climate change effects to selected food sources (based on: Tripathi et al., 2016; Sulser et al., 2021)

The climate change effect-mitigation techniques studies in the literature consist, e.g. the changes in dates of sowing, variations in the cultivars to use less water and more heat resistant crops (Kaushika et al., 2019). It is important to promote multiple cropping activities that include innovative water-saving methods and breeding approaches (Gao et al., 2019). According to the extent analysis of the correlation between grain yield and climate change, the relationship between the use of irrigation water and irrigation purposes, and human parameters such as agricultural technology innovation, policy mechanism increase and guarantee in farmland water conservancy construction financing can alleviate all types of negative effects of climate warming (Lu et al., 2019).

Many researchers even argue that the warming because of climate change may lead to a positive influence on crop production e.g. in China if concomitant alterations adopted in various cropping systems take place and positive effects of other socio-economic parameters continue in the future (Wei et al., 2014; Yang et al., 2015). Creating attention among farmers about the optimal



application of agronomic inputs under changing climatic conditions could be an effectual adaptation methodology that increases yields and may decrease their instability (Mahmood et al., 2019). Massawe et al. (2015) concluded that the main adaptation technique to minimize the effect of climate change on crop production must be through research and advancement of underutilised crops with proven potential to confront the negative effects of climate change. In such a case, the crop profile should be changed to one in which yields from other species will be satisfactory. For example, Liang et al. (2018) showed that climate change in northern China significantly affected wheat yield (+8.5% on average) but not maize yield and climate change reduced the annual nitrogen use efficiency (NUE) of maize and wheat by an average of 15%.

Adaptation of crops to climate change positively and significantly influences net crop income and productivity, so indirectly promoting the local food security for farmers, their well-being, and more advantages for people, who utilise a combination of various adaptation techniques (Abid et al., 2016). Generally, improving sustainability results need multi-scale, multi-disciplinary, multi-taxon methods to quantify the capability of ecosystems to deal with existing and future climatic factors (Alonso-Ayuso et al., 2018; Toreti et al., 2019).

### Climate change and medicinal plants

Climate changes are an important parameter affecting the distribution, growth, and reproduction of different medicinal plants (Zhang et al., 2019). Based on the predicted consequences of climate changes for many medicinal plants, their proper range will limit or move essentially (Xia et al., 2022). Rawat et al. (2022) reported that climate change will change environmental conditions in different localities and predicted the appropriate range for medicinal plants will narrow or move substantially from their original habitat. For example, *Saposhnikovia divaricata* (Turcz.) Schischk. is a notable economic medicinal plant in eastern Asia. It grows in grassy and stony slopes at 400–800 m (1,300–2,600 ft), and the margins of rice paddies, roadsides, and waste places. The range of occurrence of *S. divaricata* will increase under the influence of climate change, as the area of suitable habitats for its growth and development will increase (Chen et al., 2022). On the other hand, indicated that modifications by humans have significantly decreased the suitable distribution area of *Artemisia annua* Pall., under the current climate scenario (Wang et al., 2022).

Changes in the ranges of medicinal species concern both horizontal and vertical ranges. Climate change may not only lead to elevational shifts of mountain species but also a contraction

of their elevational ranges, which may boost the risk of local extinctions of endemic species (Zu et al., 2021). *Gentiana rigescens* Franch. is an economically important medicinal plant in the subtropical zone in Asia. It grows on grassland slopes, in valleys, in scrub, and in forests from 1100 to 3000 m above sea level. It has responded negatively to climate change. Shen et al. (2021) concluded that the areas of the current highly suitable habitat of *G. rigescens* will be negatively influenced by climate change. It is assumed that high value medicinal plants such *G. rigescens*, will face a great risk of complete extinction when the commercial requires, over-exploitation together with climate change (Applequist et al., 2020).

Extreme environmental conditions can lead to changes in the phytochemical characteristics (primary and secondary metabolites) of medicinal plants (Rawat et al., 2022). Climate change can influence the chemical composition of herbs and may induce quality changes in medicinal material (Applequist et al., 2020). Several studies have indicated that in herbal medicines the synthesis and accumulation of efficient constituents had a connection with the particular climatic environment of their habitat (Mudge et al., 2016). For example, the appropriate occurrence area of *Ziziphus jujuba* Mill. var. *spinosa* (Bunge) Hu ex H.F. Chou under the future climate scenario will be limited, as jujuboside A (triterpene saponin) and spinosin (flavone) – its active substances, are correlated with environmental parameters (Wu et al., 2022).

Climate change may boost local plant species richness, and climatic perturbation is recognised as a procedure driving biodiversity accrual. Feng et al. (2022) reported that endemic medicinal plant species richness is significantly connected with population size. Small populations are most at risk of extinction. Future climate change and human activity will therefore have a significant impact on the protection of endemic and endangered medicinal plants (Table 5).

**Tab. 5.** List of target species for highest conservation due to predicted climate changes effects (according to Cahyaningsih et al., 2021)

No	Fammily	Species name
1)	Araucariaceae	<i>Agathis borneensis</i> Warb.
2)	Apocynaceae	<i>Alstonia iwahigensis</i> Elmer
3)	Annonaceae	<i>Anaxagorea javanica</i> Blume
4)	Dipterocarpaceae	<i>Anisoptera costata</i> Pierre
5)	Thymelaeaceae	<i>Aquilaria malaccensis</i> Beneth.
6)	Acanthaceae	<i>Barleria prionitis</i> L.
7)	Fagaceae	<i>Castanopsis argentea</i> A.DC.
8)	Diksoniaceae	<i>Dicksonia blumei</i> (Kunze) T.Moore
9)	Dipterocarpaceae	<i>Dipterocarpus baudii</i> Korth.
10)	Fabaceae	<i>Euchresta horsfieldii</i> (Lesch.) Benn.
11)	Simaroubaceae	<i>Eurycoma longifolia</i> Jack

12)	Lauraceae	<i>Eusideroxylon zwageri</i> Teijsm. & Binn.
13)	Gentianaceae	<i>Gentiana quadrifaria</i> Blume
14)	Euphorbiaceae	<i>Macaranga griffithiana</i> Müll.Arg.
15)	Nepenthaceae	<i>Nepenthes reinwardtiana</i> Miq.
16)	Pinaceae	<i>Pinus merkusii</i> Jungh. & de Vriese
17)	Apocynaceae	<i>Rauvolfia serpentina</i> Benth. ex Kurz
18)	Santalaceae	<i>Santalum album</i> L.
19)	Lamiaceae	<i>Scutellaria javanica</i> Jungh.
20)	Dipterocarpaceae	<i>Shorea seminis</i> (de Vriese) Slooten

Besides, these changes could lead to modifications in plant community composition and the relative abundance of different species in the natural habitats (Getachew et al., 2021). Species with narrow ecological tolerance will be easily eliminated, and those with low habitat requirements, often with invasive features, will spread.

Probably in the future, climate changes will cause unpredictability for the reasonable introduction of medicinal plants in cultivation. It is a threat to the sustainable usage of medicinal plants. This may result in the need for greater exploitation of medicinal species from their natural habitats (Rana et al., 2020).

## Conclusions

The food system including agriculture is one of the main drivers of climate change, and climate change concurrently has significant effects on human health through the food system. An increase in temperature may lead to an increase in the chance of people being exposed to foodborne hazards, and multiple pathways are revealed to mediate climate change and increase negative health impacts through food safety. Studies on the possible impacts of climate change on healthy functional food and medicinal plants are particularly significant because of their value within traditional systems of medicine and as economically useful plants.

[1] In general, when plants are subjected to environmental stress, the production of secondary metabolites may increase because somatic growth is often inhibited more than photosynthesis, and growth-related carbon is instead transferred to secondary metabolites. Different plant species will react individually to climate change. Some species will remain in the region but adapt to new climatic conditions via selection, other species will shift to higher altitudes or latitudes, and some species of medicinal plants may become extinct.

[2] During the growing season, important phenological events for medicinal and healthy functional plants are adapted to the given climate. These include flowering and fruit setting, bud bursting and

leaf development, and leaf fall in autumn or during the dry season. However, global warming processes have and will have an impact in the future on the timing of spring and the length of the growing season. This determines the course of plant phenological processes, affecting their productivity.

[3] Climate change impacts on overall global food production are likely destructive, especially later in the century, but these impacts vary mainly by region. The most negative impacts are possibly to occur in the subtropics and tropics. These negative effects can be mitigated by implementing adaptation programs based on solid scientific principles and taking into account local biophysical and socio-economic conditions. In addition, further research and monitoring is necessary to assess the impact of climate change on the yields and chemical components of medicinal plants and health-promoting functional foods.

#### Conflict of interest

The author declare no conflict of interest related to this article.

#### References

- Abid, M., Schneider, U.A., Scheffran, J. (2016). Adaptation to climate change and its impacts on food productivity and crop income: Perspectives of farmers in rural Pakistan. *Journal of Rural Studies*, 47, 254–266. <https://doi.org/10.1016/j.jrurstud.2016.08.005>
- Alonso-Ayuso, M., Quemada, M., Vanclooster, M., Ruiz-Ramos, M., Rodriguez, A., Gabriel, J.L. (2018). Assessing cover crop management under actual and climate change conditions. *Science of Total Environment*, 621, 1330–1341. <https://doi.org/10.1016/j.scitotenv.2017.10.095>
- Alvi, S., Roson, R., Sartori, M., Jamil, F. (2021). An integrated assessment model for food security under climate change for South Asia. *Heliyon*, 7(4), e06707. <https://doi.org/10.1016/j.heliyon.2021.e06707>
- Applequist, W.L., Brinckmann, J.A., Cunningham, A.B., Hart, R.E., Heinrich, M., Katerere, D.R., Andel, T.V. (2020). Scientists' warning on climate change and medicinal plants. *Planta Medica*, 86(1), 10–18. <https://doi.org/10.1055/a-1041-3406>
- Auerswald, K., Menzel, A. (2021). Change in erosion potential of crops due to climate change. *Agricultural and Forest Meteorology*, 300, 108338. <https://doi.org/10.1016/j.agrformet.2021.108338>
- Ayllon, D., Railsback, S.F., Harvey, B.C., Quiros, I.G., Nicola, G.G., Elvira, B., Almodovar, A. (2019). Mechanistic simulations predict that thermal and hydrological effects of climate change on Mediterranean trout can not be offset by adaptive behaviour, evolution, and increased food production. *Science of Total Environment*, 693, 133648. <https://doi.org/10.1016/j.scitotenv.2019.133648>
- Bates, A.E., Cooke, R.S.C., Duncan, M.I., Edgar, G.J., Bruno, J.F., Benedetti-Cecchi, L., Cote, I.M., Lefcheck, J.S., Costello, M.J., Barrett, N., Bird, T.J., Fenberg, P.B., Stuart-Smith, R.D. (2019). Climate resilience in marine

- protected areas and the Protection Paradox. *Biological Conservation*, 236, 305–314. <https://doi.org/10.1016/j.biocon.2019.05.005>
- Bayir, G.B, Aksoy, A.N., Kocyigit, A. 2019. The Importance of Polyphenols as Functional Food in Health. *Bezmi'alem Science*, 7(2), 157–163. <https://doi.org/10.14235/bas.galenos.2018.2486>
- Butnariu, M., Sarac, I. (2019). Functional food. *International Journal of Nutrition*, 3(3), 7–16. <https://doi.org/10.14302/issn.2379-7835.ijn-19-2615>
- Cahyaningsih, R., Phillips, J., Brehm, J.M., Gaisberger, H., Maxted, N. (2021). Climate change impact on medicinal plants in Indonesia. *Global Ecology and Conservation*, 30, e01752. <https://doi.org/10.1016/j.gecco.2021.e01752>
- Cencis, A., Chingwaru, W. (2010). The role of functional foods, nutraceuticals, and food supplements in intestinal health. *Nutrient*, 2, 611–625. <https://doi.org/10.3390/nu2060611>
- Chen, B., Zou, H., Zhang, B., Zhang, X., Jin, X., Wang, C., Zhang, X. (2022). Distribution pattern and change prediction of *Saposhnikovia divaricata* suitable area in China under climate change. *Ecological Indicators*, 143, 109311. <https://doi.org/10.1016/j.ecolind.2022.109311>
- Choudhary, R., Tandon, R.V. (2009). Consumption of functional food and our health concerns. *Pakistan Journal of Physiology*, 5(1), 76–83.
- Cobbinah, P.B., Asibey, M.O., Opoku-Gyamfi, M., Peprah, C. (2019). Urban planning and climate change in Ghana. *Journal of Urban Managing*, 8, 261–271. <https://doi.org/10.1016/j.jum.2019.02.002>
- Das, R., Biswas, S., Banerjee, E.R. (2016). Nutraceutical-prophylactic and therapeutic role of functional food in health. *Journal of Nutrition and Food Sciences*, 6(4), 1–17. <https://doi.org/10.4172/2155-9600.1000527>
- Del Castillo, M.D., Iriundo-DeHond, A., Martirosyan, D.M. (2018). Are functional foods essential for sustainable health? *Annals of Nutrition and Food Science*, 2(1), 1015. <https://doi.org/10.3390/nu10101358>
- Dorji, T., Hopping, K.A., Meng, F., Wang, S., Jiang, L., Klein, J.A. (2020). Impacts of climate change on flowering phenology and production in alpine plants: The importance of end of flowering. *Agriculture Ecosystem and Environment*, 291, 106795. <https://doi.org/10.1016/j.agee.2019.106795>
- Duchenne-Moutien, R.A., Neetoo, H. (2021). Climate change and emerging food safety issues: A review. *Journal of Food Protection*, 84(11), 1884–1897. <https://doi.org/10.4315/JFP-21-141>
- Feng, G., Xiong, Y.-J., Wei, H.-Y., Li, Y., Mao, L.-F. (2022). Endemic medicinal plant distribution correlated with stable climate, precipitation, and cultural diversity. *Plant Diversity*, 45(4), 479–484. <https://doi.org/10.1016/j.pld.2022.09.007>
- Gao, J., Yang, X., Zheng, B., Liu, Z., Zhao, J., Sun, S., Li, K., Dong, C. (2019). Effects of climate change on the extension of the potential double cropping region and crop water requirements in Northern China. *Agricultural and Forest Meteorology*, 268, 146–155. <https://doi.org/10.1016/j.agrformet.2019.01.009>
- Getachew, F., Bayabil, H.K., Hoogenboom, G., Teshome, F.T., Zewdu, E. (2021). Irrigation and shifting planting date as climate change adaptation strategies for sorghum. *Agriculture Water Management*, 255, 106988. <https://doi.org/10.1016/j.agwat.2021.106988>
- Gohari, A., Eslamian, S., Abedi-Koupaei, J., Bavani, A.M., Wang, D., Madani, K. (2013). Climate change impacts on crop production in Irans Zayandeh-Rud river basin. *Science of Total Environment*, 442, 405–419. <https://doi.org/10.1016/j.scitotenv.2021.10.029>

- Green, R., Scheelbeek, P., Bentham, J., Cuevas, S., Smith, P.P., Dangour, P.A. (2022). Growing health: global linkages between patterns of food supply, sustainability, and vulnerability to climate change. *Lancet Planetary Health*, 6(11), 901–908. [https://doi.org/10.1016/S2542-5196\(22\)00223-6](https://doi.org/10.1016/S2542-5196(22)00223-6)
- Guo, H., Huang, Z., Tan, M., Tuan, H., Awe, G.O., Are, K.S., Abegnrin, T.P., Hussain, Z., Kuang, Z., Liu, D. (2021). Crop resilience to climate change: A study of spatio-temporal variability of sugarcane yield in a subtropical region, China. *Smart Agricultural Technology*, 1, 100014. <https://doi.org/10.1016/j.atech.2021.100014>
- Kang, X., Qi, J., Li, S., Meng, F.-R. (2022). A watershed-scale assessment of climate change impacts on crop yields in Atlantic Canada. *Agriculture and Water Management*, 269, 107680. <https://doi.org/10.1016/j.agwat.2022.107680>
- Katsini, L., Vhonsale, S., Akkermans, S., Roufou, S., Griffin, S., Valdramidis, V., Misiou, O., Koutsoumanis, K., Lopez, C.A.M., Polanska, M., Impe, J.F.M.V. (2022). Quantitative methods to predict the effect of climate change on microbial food safety: A needs analysis. *Trends in Food Science and Technology*, 126, 113–125. <https://doi.org/10.1016/j.tifs.2021.07.041>
- Kaushika, G.S., Arora, H., Prasad, H. (2019). Analysis of climate change effects on crop water availability for paddy, wheat and berseem. *Agriculture and Water Management*, 225, 105734. <https://doi.org/10.1016/j.agwat.2019.105734>
- Kotilainen, L., Rajalahti, R., Ragasa, C., and Pehu, E. (2006). Health enhancing foods: Opportunities for strengthening the sector in developing countries. *Agriculture and Rural Development discussion paper no. 30*, Washington, D.C.: World Bank Group. <http://documents.worldbank.org/curated/en/314421468324275640/Health-enhancing-foods-opportunities-for-strengthening-developing-countries>
- Lemes, P., Barbosa, F.G., Naimi, B., Araujo, M.B. (2022). Dispersal abilities favor commensalism in animal-plant interactions under climate change. *Science of Total Environment*, 835, 155157. <https://doi.org/10.1016/j.scitotenv.2022.155157>
- Li, Y., Tian, D., Feng, G., Yang, W., Feng, L. (2021). Climate change and cover crop effects on water use efficiency of a corn-soybean rotation system. *Agriculture and Water Management*, 255, 107042. <https://doi.org/10.1016/j.agwat.2021.107042>
- Liang, S., Li, Y., Zhang, X., Sun, Z., Sun, N., Duan, Y., Xu, M., Wu, L. (2018). Response of crop yield and nitrogen use efficiency for wheat-maize cropping system to future climate change in northern China. *Agricultural and Forest Meteorology*, 262, 310–321. <https://doi.org/10.1016/j.agrformet.2018.07.019>
- Lu, S., Bai, X., Li, W., Wang, N. (2019). Impacts of climate change on water resources and grain production. *Technol Forecast Social Change*, 143, 76–84. <https://doi.org/10.1016/j.techfore.2019.01.015>
- Mahmood, N., Arshad, M., Kachele, H., Ma, H., Ullah, A., Muller, K. (2019). Wheat yield response to input and socioeconomic factors under changing climate: Evidence from rainfed environments of Pakistan. *Science of Total Environment*, 688, 1275–1285. <https://doi.org/10.1016/j.scitotenv.2019.06.266>
- Mahmoudpour, Z., Shirafkan, H., Mojahedi, M., Gorji, N., Mozaffarpur, S.A. (2018). Digesters in traditional Persian medicine. *Caspian Journal of Internal Medicine*, 9(1), 1–6. <https://doi.org/10.18502/tim.v7i3.10766>
- Massawe, F.J., Mayes, S., Cheng, A., Chai, H.H., Cleasby, P., Symonds, R., Ho, W.K., Siise, A., Wong, Q.N., Kendabie, P., Yanusa, Y., Jamalluddin, N., Singh, A., Azman, R., Azam-Ali, S.N. (2015). The potential for undertutilised crops to improve food security in the face of climate change. *Procedia Environmental Sciences*, 29, 140–141.

<https://doi.org/10.1016/j.proenv.2015.07.228>

- Miron, I.J., Linares, C., Diaz, J. (2023). The influence of climate change on food production and food safety. *Environmental Research*, 216(3), 114674. <https://doi.org/10.1016/j.envres.2022.114674>
- Mudge, E., Applequist, W.L., Finley, J., Lister, P., Townesmith, A.K., Walker, K.M., Brown, P.N. (2016). Variation of select flavonols and chlorogenic acid content of elderberry collected throughout the Eastern United States. *Journal of Food Composition and Analysis*, 47, 2–59. <https://doi.org/10.1016/j.jfca.2015.12.003>
- Newbery, F., Qi, A., Fitt, B.D.L. (2016). Modelling impacts of climate change on arable crop diseases: progress, challenges and applications. *Current Opinion in Plant Biology*, 32, 101–109. <https://doi.org/10.1016/j.pbi.2016.07.002>
- Phillips, J.D. (2019). State factor network analysis of ecosystem response to climate change. *Ecological Complexity*, 40, 100789. <https://doi.org/10.1016/j.ecocom.2019.100789>
- Ponce, C. (2020). Intra-seasonal climate variability and crop diversification strategies in the Peruvian Andes: A word of caution on the sustainability of adaptation to climate change. *World Development*, 127, 104740. <https://doi.org/10.1016/j.worlddev.2019.104740>
- Raes, D., Waongo, M., Vanuytrecht, E., Moreno, P.M. (2021). Improved management alleviate some but not all of the adverse effects of climate change on crop yields in smallholder farms in West Africa. *Agricultural and Forest Meteorology*, 309, 108563. <https://doi.org/10.1016/j.agrformet.2021.108563>
- Raiten, D.J., Bremer, A.A. (2023). Exploring the intersection of climate/environmental change, food systems, nutrition, and health: global challenge, opportunity, or both? *American Journal of Clinical Nutrition*, 117(2), 224–226. <https://doi.org/10.1016/j.ajcnut.2022.11.024>
- Ranga Rao, A., Ravishankar, G.A. (2018). Algae as source of functional ingredients for health benefits. *Agricultural Research and Technology Open Access Journal*, 14(2), 555911. <https://doi.org/10.19080/artoaj.2018.14.555911>
- Rawat, N., Purohit, S., Painuly, V., Negi, G.S., Bisht, M.P.S. (2022). Habitat distribution modeling of endangered medicinal plant *Picrorhiza kurroa* (Royle ex Benth) under climate change scenarios in Uttarakhand Himalaya, India. *Ecological Informatics*, 68, 101550. <https://doi.org/10.1016/j.ecoinf.2021.101550>
- Raza, M.M., Bebbler, D.P. (2022). Climate change and plant pathogens. *Current Opinion in Microbiology*, 70, 102233. <https://doi.org/10.1016/j.mib.2022.102233>
- Rolo, V., Moreno, G. (2019). Shrub encroachment and climate change increase the exposure to drought of Mediterranean wood-pastures. *Science of the Total Environment*, 660, 550–558. <https://doi.org/10.1016/j.scitotenv.2019.01.029>
- Ruszkiewicz, J.A., Tinkov, A.A., Skalny, A.V., Siokas, V., Dardiotis, E., Tsatsakis, A., Bowman, A.B., da Rocha, J.B.T., Aschner, M. (2019). Brain diseases in changing climate. *Environmental Research*, 177, 108637. <https://doi.org/10.1016/j.envres.2019.108637>
- Seidel, S.J., Gaiser, T., Ahrends, H.E., Hugging, H., Siebert, S., Bauke, S.L., Gocke, M.I., Koch, M., Schweitzer, K., Schaaf, G., Ewert, F. (2021). Crop response to P fertilizer omission under a changing climate- Experimental and modeling results over 115 years of a long-term fertilizer experiment. *Field Crops Research*, 268, 108174. <https://doi.org/10.1016/j.fcr.2021.108174>
- Semba, R.D., Askari, S., Gibson, S., Bloem, M.W., Kraemer, K. (2022). The potential impact of climate change on the

- micronutrient-rich food supply. *Advances in Nutrition*, 13(1), 80–100. <https://doi.org/10.1093/advances/nmab104>
- Shahzad, Z., Rouached, H. (2022). Protecting plant nutrition from the effects of climate change. *Current Biology*, 32(13), E725–R727. <https://doi.org/10.1016/j.cub.2022.05.056>
- Shandilya, U.K., Sharma, A. (2017). Functional foods and their benefits: an overview. *Journal of Nutritional Health and Food Engineering*, 7(4), 353–356. <https://doi.org/10.15406/jnhfe.2017.07.00247>
- Sharma, H., Rao, P.S., Singh, A.K. (2021). Fifty years of research on *Tinospora cordifolia*: from botanical plant to functional ingredient in foods. *Trends in Food Science and Technology, Part A*, 118, 189–206. <https://doi.org/10.1016/j.tifs.2021.10.003>
- Shen, T., Yu, H., Wang, Y.-Z. (2021). Assessing the impacts of climate change and habitat suitability on the distribution and quality of medicinal plant using multiple information integration: Take *Gentiana rigescens* as an example. *Ecological Indicators*, 123, 107376. <https://doi.org/10.1016/j.ecolind.2021.107376>
- Sulser, T.B., Beach, R.H., Wiebe, K., Dunston, S., Fukagawa, N.K. (2021). Disability-adjusted life years due to chronic and hidden hunger under food system evolution with climate change and adaptation to 2050. *The American Journal of Clinical Nutrition*, 114(2), 550–563. <https://doi.org/10.1093/ajcn/nqab101>
- Sun, W., Shahrajabian, M.H., Petropoulos, S.A., Shahrajabian, N. (2023). Developing sustainable agriculture systems in medicinal and aromatic plant production by using chitosan and chitin-based biostimulants. *Plants*, 12(13), 2469. <https://doi.org/10.3390/plants12132469>
- Sun, W., Shahrajabian, M.H., Soleymani, A. (2024). The roles of plant-growth-promoting rhizobacteria (PGPR)-based biostimulants for agricultural production systems. *Plants*, 13(5), 613. <https://doi.org/10.3390/plants13050613>
- Supachaturat, S., Pichyangkura, R., Chandrachai, A., Pentrakoon, D. (2017). Perspective on functional food commercialization in Thailand. *International Food Research Journal*, 24(4), 1374–1382. <https://doi.org/10.7763/ijssh.2015.v5.560>
- Teixeira, E., Kersebaum, K.C., Ausseil, A.-G., Cichota, R., Guo, J., Johnstone, P., George, M., Liu, J., Malcolm, B., Khaembah, E., Meiyalaghan, S., Richards, K., Zyskowski, R., Michel, A., Sood, A., Tait, A., Ewert, F. (2021). Understanding spatial and temporal variability of N leaching reduction by winter cover crops under climate change. *Science of Total Environment*, 771, 144770. <https://doi.org/10.1016/j.scitotenv.2020.144770>
- Toreti, A., Bassu, S., Ceglar, A., Zampieri, M. (2019). Climate change and crop yields. *Encyclopedia of Food Security and Sustainability*, 1, 223–227. <https://doi.org/10.1016/B978-0-08-100596-5.22010-6>
- Tripathi, A., Tripathi, D.K., Chauhan, D.K., Kumar, N., Singh, G.S. (2016). Paradigms of climate change impacts on some major food sources of the world: A review on current knowledge and future prospects. *Agriculture, Ecosystem, Environment*, 216, 356–373. <https://doi.org/10.1016/j.agee.2015.09.034>
- Wang, D., Shi, C., Alamgir, K., Kwon, S.M., Pan, L., Zhu, Y., Yang, X. (2022). Global assessment of the distribution and conservation status of a key medicinal plant (*Artemisia annua* L.): The roles of climate and anthropogenic activities. *Science of Total Environment*, 821, 153378. <https://doi.org/10.1016/j.scitotenv.2022.153378>
- Wang, Y., Liu, S., Shi, H. (2023). Comparison of climate change impacts on the growth of C3 and C4 crops in China. *Ecological Informatics*, 74, 101968. <https://doi.org/10.1016/j.ecoinf.2022.101968>
- Weber, L., Bartek, L., Brancoli, P., Sjolund, A., Eriksson, M. (2023). Climate change impact of food distribution: The case of reverse logistics for bread in Sweden. *Sustainable Production and Consumption*, 36, 386–396.



<https://doi.org/10.1016/j.spc.2023.01.018>

- Wei, T., Cherry, T.L., Glomrod, S., Zhang, T. (2014). Climate change impacts on crop yield: Evidence from China. *Science of Total Environment*, 499, 133–140. <https://doi.org/10.1016/j.scitotenv.2014.08.035>
- Wu, M., Gu, X., Zhang, Z., Si, M., Zhang, Y., Tian, W., Ma, D. (2022). The effects of climate change on the quality of *Ziziphus jujuba* var. *Spinosa* in China. *Ecological Indicators*, 139, 108934. <https://doi.org/10.1016/j.ecolind.2022.108934>
- Xia, C., Huang, Y., Qi, Y., Yang, X., Xue, T., Hu, R., Deng, H., Bussmann, R.W., Yu, S. (2022). Developing long-term conservation priority planning for medicinal plants in China by combining conservation status with diversity hotspot analyses and climate change prediction. *BMC Biology*, 20(89), 1–20. <https://doi.org/10.1186/s12915-022-01285-4>
- Yang, X., Chen, F., Lin, X., Liu, Z., Zhang, H., Zhao, J., Li, K., Ye, Q., Li, Y., Lv, S., Yang, P., Wu, W., Li, Z., Lal, R., Tang, H. (2015). Potential benefits of climate change for crop productivity in China. *Agricultural and Forest Meteorology*, 208, 76–86. <https://doi.org/10.1016/j.agrformet.2015.04.024>
- Yeung, A.W.K., Mocan, A., Atanasov, A.G. (2018). Let food be thy medicine and medicine be thy food: A bibliometric analysis of the most cited papers focusing on nutraceuticals and functional foods. *Food Chemistry*, 269, 455–465. <https://doi.org/10.1016/j.foodchem.2018.06.139>
- Yin, R., Gruss, I., Eisenhauer, N., Kardol, P., Thakur, M.P., Schmidt, A., Xu, Z., Siebert, J., Zhang, C., Wu, G.-L., Schadler, M. (2019). Land use modulates the effects of climate change on density but not community composition of Collembola. *Soil Biology and Biochemistry*, 138, 107598. <https://doi.org/10.1016/j.soilbio.2019.107598>
- Zhang, J., Liu, Y. (2022). Decoupling of impact factors reveals the response of cash crops phenology to climate change and adaptive management practice. *Agricultural and Forest Meteorology*, 322, 109010. <https://doi.org/10.1016/j.agrformet.2022.109010>
- Zhang, K., Zhang, Y., Zhou, C., Meng, J., Sun, J., Zhou, T., Tao, J. (2019). Impact of climate factors on future distributions of *Paeonia ostii* across China estimated by MaxEnt. *Ecological Informatics*, 50, 62–67. <https://doi.org/10.1016/j.ecoinf.2019.01.004>
- Zhang, Z., Li, Y., Chen, X., Wang, Y., Niu, B., Liu, D.L., He, J., Pulatov, B., Hassan, I., Meng, Q. (2023). Impact of climate change and planting date shifts on growth and yields of double cropping rice in southeastern China in future. *Agricultural Systems*, 205, 103581. <https://doi.org/10.1016/j.agsy.2022.103581>
- Zu, K., Wang, Z., Zhu, X., Lenoir, J., Shrestha, N., Lyu, T., Luo, A., Li, Y., Ji, C., Peng, S., Meng, J., Zhou, J. (2021). Upward shift and elevational range contractions of subtropical mountain plants in response to climate change. *Science of Total Environment*, 783, 146896. <https://doi.org/10.1016/j.scitotenv.2021.146896>

# Wpływ zmian klimatycznych na zdrową żywność funkcjonalną oraz rośliny lecznicze

## Streszczenie

Istnieje coraz większa potrzeba badań mających na celu określenie wpływu zmian klimatycznych na fenologię roślin leczniczych i prozdrowotnych, przemian ich siedlisk, zmian zasięgu oraz produkcję metabolitów wtórnych. Promowanie świadomości na temat potencjalnych skutków ocieplenia (m.in. podwyższonego stężenia CO<sub>2</sub> i zwiększonego promieniowania ultrafioletowego), wpływających na składniki wtórne roślin i metabolity, jest ważnym zadaniem na przyszłość. Istnieją dowody, że zmiany klimatyczne powodują następstwa objawiające się zmianami cyklu życia i rozmieszczenia roślin, zarówno tych stanowiących żywność funkcjonalną, jak i dzikorosnących taksonów leczniczych. Zmuszają one rośliny do aklimatyzacji i zmian w fenologii oraz do rozwoju nowych cech fizycznych. Szybko zmieniający się klimat może być korzystny dla gatunków, które mogą szybko rozszerzyć swoje zasięgi lub które mogą tolerować szeroki zakres warunków klimatycznych, a obie te cechy są wspólne dla różnych gatunków roślin inwazyjnych. Niniejszy przegląd zawiera podsumowanie wpływu zmian klimatycznych na rośliny lecznicze, prozdrowotne i stanowiące żywność funkcjonalną.

**Słowa kluczowe:** czynniki klimatyczne, globalne ocieplenie, plony roślin, zasięgi gatunków

## Information on the author

**Mohamad Hesam Shahrajabian** <https://orcid.org/0000-0002-8638-1312>

His research interests focus on broadly understood issues related to agronomy and plant breeding and yielding. He cooperates with, among others, the Biotechnology Research Institute, Chinese Academy of Agricultural Sciences in Beijing.